MODULATOR OF INTRACELLULAR Ca^{2+} , THAPSIGARGIN, INTERFERES WITH *IN VITRO* SECRETION OF CYTOKINES AND NITRIC OXIDE

Eva Kmoníčková^{a,b}, Nikolina Kutinová Canová^b, Hassan Farghali^b, Antonín Holý^c,
Zdeněk Zidek^a

^a Institute of Experimental Medicine, Academy of Sciences, 142 20 Prague, Czech Republic

^b Institute of Pharmacology, 1st Faculty of Medicine, Charles University, 128 00 Prague, Czech Republic

^c Institute of Organic Chemistry and Biochemistry, Academy of Sciences, 166 00 Prague, Czech Republic
e-mail: kmonickova@biomed.cas.cz

Received: June 10, 2005; Accepted: September 25, 2005

Key words: Thapsigargin/TNF- α /IL-10/RANTES/MIP-1 α

Interference of thapsigargin (TG), an inhibitor of endoplasmic reticulum Ca^{2+} ATPase, with immune reactivity of murine macrophages was investigated under conditions *in vitro*. The activation of cells with lipopolysaccharide (LPS), interferon- γ (IFN- γ), and with acyclic nucleoside phosphonate *N*⁶-isobutyl-9-[2-(phosphonomethoxy)ethyl]-2,6-diaminopurine (*N*⁶-isobutyl-PMEDAP) resulted in enhanced production of cytokines TNF- α , IL-10, chemokines RANTES/CCL5 and MIP-1 α /CCL3, as well as in substantially augmented production of nitric oxide (NO) triggered by IFN- γ . The effects were in a dual mode of action influenced by TG (1 μM). While TG upregulated secretion of TNF- α , it inhibited secretion of IL-10 and RANTES. The immune-stimulated secretion of MIP-1 α remained virtually unaffected, though TG on its own activated expression of MIP-1 α in macrophages. The high-output NO production induced by IFN- γ , high concentrations of LPS, or by combination of IFN- γ plus LPS or *N*⁶-isobutyl-PMEDAP was inhibited by TG. On the other hand, production of NO which was marginally activated by low concentration of LPS was upregulated by TG.

INTRODUCTION

Physiologically maintained level of cytosolic free Ca^{2+} is a crucial signal controlling many cell functions^{1,2}, and any imbalance may have cytotoxic consequences³. Also the role of calcium ions in the expression of immunomodulatory properties of drugs and their importance in effectiveness of natural defence mechanisms are being intensively explored. A plausible experimental approach to study this problem is the use of modulators of intracellular calcium. Thapsigargin (TG) is a naturally occurring sesquiterpene lactone with prominent acute action on various types of cells. It is widely employed as a selective and irreversible inhibitor of endoplasmic reticulum Ca^{2+} ATPase leading to rapid elevation of intracellular concentration of free calcium. The interference of TG with immune functions of cells is largely unknown. We investigated its effects on production of cytokines including chemokines RANTES and MIP-1 α under conditions *in vitro*.

MATERIALS AND METHODS

Animals and isolation of peritoneal macrophages

Female mice of the inbred strain C57BL/6, 7–9 wks old, were purchased from Charles River Deutschland (Sulzfeld, Germany). They were kept in transparent plastic cages in groups of eight, and maintained in an

Independent Environmental Air Flow Cabinet (ESI Flufrance, Wissous, France). Lighting was set on 0600 to 1800 h, temperature at 22 °C.

Mice, sacrificed by cervical dislocation, were injected intraperitoneally with 8 ml sterile saline. Collected and pooled lavage cells were washed, resuspended in culture medium and seeded into 96-well flat-bottom Nunc microplates in 100- μl volumes. Adherent peritoneal cells (macrophages) were isolated by incubating the cells for 2 h at 37 °C, 5% CO_2 , and then thrice vigorously shaking the plate and washing to remove non-adherent cells. They were cultured with or without addition of immunostimulatory agents, i.e. interferon- γ (IFN- γ ; 5000 pg/ml), lipopolysaccharide (LPS; 10–10.000 pg/ml), and *N*⁶-isobutyl-9-[2-(phosphonomethoxy)ethyl]-2,6-diaminopurine (*N*⁶-isobutyl-PMEDAP; 2–100 μM), and in the absence or presence of thapsigargin (TG; 1 μM) in Heraeus incubator for 24 h. TG was added 30 min before the immunostimulators. All experimental variants were set in duplicate.

Culture medium and chemicals

Complete RPMI-1640 medium contained 10 % heat-inactivated fetal bovine serum, 2 mM L-glutamine, 50 $\mu\text{g}/\text{ml}$ gentamicin, and 5×10^{-5} M 2-mercaptoethanol (all Sigma). Recombinant mouse IFN- γ was purchased from R&D Systems (Minneapolis, MN), LPS of *S. typhimurium* origin from Difco Labs (Detroit, MI). Acyclic nucleoside phosphonate *N*⁶-isobutyl-9-[2-

Fig. 1. Effect of thapsigargin (TG, 1 μ M) on 5-h secretion of cytokines by mouse macrophages. The following concentrations of primary immune stimulators were applied: LPS 1 ng/ml, IFN- γ 5 ng/ml, *N*⁶-isobutyl-PMEDAP (PMEDAP) 50 μ M. The data are means $\pm$ SEM.

(phosphonomethoxy)ethyl]-2,6-diaminopurine (*N*⁶-isobutyl-PMEDAP) was synthesized in-house (Inst. Organic Chemistry and Biochemistry) according to the procedure described elsewhere⁴. The cell viability and cell proliferation assay kit WST-1 was bought from Roche Applied Science (Mannheim, Germany). Thapsigargin was obtained from Sigma-Aldrich (Prague, CZ).

Immunobiological assays

Concentration of cytokines and chemokines in the 5-h cell supernatants was determined using enzyme-linked immunoabsorbent assay (ELISA) kits, following the manufacturer's instructions (R&D Systems, Minneapolis, MN).

The concentration of nitrites in the 24-h supernatants was taken as a measure of NO production. It was detected in individual cell-free samples (50 μ l) incubated for 10 min at 37 $^{\circ}$ C with an aliquot of a Griess reagent (1% sulphanilamide/0.1% naphthylendiamine/2.5% H_3PO_4). The absorbance at 540 nm was recorded using a microplate

spectrophotometer (Tecan, Austria). A nitrite calibration curve was used to convert absorbance to μ M nitrite.

Viability of macrophages was assayed after the 20 h of culture, following the manufacturer's instructions (Roche). Absorbances at 450 nm were determined at the interval of 2.5 h after addition of WST-1.

Data analysis

Analysis of variance (ANOVA) with subsequent Bonferroni's multiple comparison test for selected pairs, and graphical presentation of data were done using the Prism program (GraphPad Software, San Diego, CA).

RESULTS AND DISCUSSION

Irrespective of the type of primary immune stimulation of macrophages, TG significantly augmented secretion of proinflammatory cytokine TNF- α (Fig. 1). This is in good line with the observation that transient increase

Fig. 2. Interference of thapsigargin (TG) with production of NO by mouse macrophages. NO was activated by IFN- γ , LPS, *N*⁶-isobutyl-PMEDAP (PMEDAP) and their combinations. Thapsigargin was added 30 min before. Supernatant concentration of nitrites was determined after the 24-h culture. Each point is mean $\pm$ SEM and data are representative of two identical experiments.

of intracellular calcium plays a role in the LPS-induced expression of TNF- α in macrophage cell line J774.1 (ref.⁵). However, an opposite effect was observed in hepatocytes where TG, in dependence on experimental conditions, did not block or rather inhibited release of TNF- α (ref.⁶). In accordance with the finding of immunostimulatory potential of TG are data showing an enhancement of IL-8 production by human colonic epithelial cells⁷, and IL-6 by rat⁸, and mouse⁹ peritoneal macrophages. In addition, the release of calcium from endoplasmic reticulum stores has been found to activate secretion of IL-1 α and IL-1 β in murine macrophages¹⁰. We found that TG had virtually no effect on the immune-stimulated secretion of chemokine MIP-1 α /CCL3, although it did activate secretion of the chemokine in immune-nonstimulated macrophages. In sharp contrast, TG profoundly inhibited production of chemokine RANTES/CCL5 as well as antiinflammatory cytokine IL-10 (Fig. 1).

Although secretion of the major NO-upregulatory cytokine TNF- α was enhanced, production of NO was deeply suppressed by TG (Fig. 2) in combination with IFN- γ , IFN- γ /LPS, PMEDAP, IFN- γ /PMEDAP, and with relatively high dose of LPS (10 ng/ml). Notably, as low concentration of TG as 0.1 μ M was already effective. On the other hand, production of NO following the low dose of LPS (10 pg/ml) was significantly potentiated by TG. The data are in agreement with findings demonstrating both NO-stimulatory¹¹ and NO-inhibitory¹² effects of

Fig. 3. Effect of thapsigargin on formation of formazan from the tetrazolium salts by mouse macrophages. The WST-1 assay was done after the 20-h culture of cells in the presence of IFN- γ , LPS, and *N*⁶-isobutyl-PMEDAP (PMEDAP; 50 μ M) $\pm$ thapsigargin (1 μ M). Each bar represents mean $\pm$ SEM obtained from six culture wells.

TG on iNOS mRNA expression and NO biosynthesis in murine macrophages.

We have found that TG decreases formation of formazan from the tetrazolium salts (Fig. 3), suggesting its toxic influence in terms of decreased number of metabolically active cells in the culture. The suppression was very similar in all experimental settings, reaching statistically highly significant over-all value of 30% ($F_{1,6} = 250$; $P < 0.0001$). With respect to the fact that not only inhibitory but also stimulatory effects of TG have been observed, the cytotoxicity of TG is unlikely to explain its immunomodulatory interventions.

It can be concluded that TG interferes with cell immune functions. Our original data show that it also interferes with expression of chemokines. The antithetical effects suggest that TG may target multiple underlying signaling pathways. Toxicological aspects of the dual mode of action remain to be firmly established.

ACKNOWLEDGEMENT

This study was supported by grants from the Grant agency of the Czech Republic # 305/05/2425, # 305/03/1470, and from the Centre for New Antivirals and Antineoplastics IM6138896301. It was performed as a part of research project of the Institute of Experimental Medicine # AVOZ50390512.

REFERENCES

1. Berridge MJ. (1997) Elementary and global aspects of calcium signalling. *J Physiol (London)* 499, 291–306.
2. Bootman MD, Berridge MJ, Lipp P. (1993) Cooking with calcium: the recipes for composing global signals from elementary events. *Cell* 91, 367–73.
3. Annunziato L, Amoroso S, Pannaccione A, Cataldi M, Pignataro G, D'Alessio A, Sirabella R, Secondo A, Sibaud L, Di Renzo GF. (2003) Apoptosis induced in neuronal cells by oxidative stress: role played by caspases and intracellular calcium ions. *Toxicol Lett* 139, 125–33.
4. Holý A, Votruba I, Tloušťová E, Masojídková M. (2001) Synthesis and cytostatic activity of *N*-[2-(phosphonomethoxy)alkyl] derivatives of *N*⁶-substituted adenines, 2,6-diaminopurines and related compounds. *Collect Czech Chem Commun* 66, 1545–92.
5. Watanabe N, Suzuki J, Kobayashi Y. (1996) Role of calcium in tumor necrosis factor- α production by activated macrophages. *J Biochem* 120, 1190–95.
6. Lichtman SN, Wang J, Zhang C, Lemasters JJ. (1996) Endocytosis and Ca^{2+} are required for endotoxin-stimulated TNF- α release by rat Kupffer cells. *Am J Physiol* 271, G920–8.
7. Yu Y, De Waele C, Chadee K. (2001) Calcium-dependent interleukin-8 gene expression in T84 human colonic epithelial cells. *Inflamm Res* 50, 220–6.
8. Ichinowatari G, Yamada M, Yaginuma H, Tsuyuki K, Tanimoto A, Ohuchi, K. (2002) Participation of prostaglandin E_2 and platelet-activating factor in thapsigargin-induced production of interleukin-6. *Eur J Pharmacol* 434, 187–96.
9. Bost KL, Mason MJ. (1995) Thapsigargin and cyclopiazonic acid initiate rapid and dramatic increases of IL-6 mRNA expression and IL-6 secretion in murine peritoneal macrophages. *J Immunol* 155, 285–96.
10. Brough D, Le Feuvre RA, Wheeler RD, Solovyova N, Hilfiker S, Rothwell NJ, Verkhatsky A. (2003) Ca^{2+} stores and Ca^{2+} entry differentially contribute to the release of IL-1 β and IL-1 α from murine macrophages. *J Immunol* 170, 3029–36.
11. Park YC, Jun CD, Kang HS, Kim HD, Chung HT. (1995) Intracellular Ca^{2+} pool depletion is linked to the induction of nitric oxide synthesis in murine peritoneal macrophages. *Biochem Mol Biol Int* 36, 949–55.
12. Jordan ML, Rominski B, Jaquins-Gerstl A, Geller D, Hoffman RA. (1995) Regulation of inducible nitric oxide production by intracellular calcium. *Surgery* 118, 138–45.