

The application of volumetry as an indication criterion in blow-out fractures

Daniel Kovar^{a,b}, Richard Holy^a, Zdenek Voldrich^a, Pavel Voska^a, Jan Lestak^c, Jaromir Astl^a

Background and Aim. This work builds on our publication on the subject of creating a mathematical model for calculating the volume of prolapsed soft tissue of the orbit in blow-out fractures, which aids us greatly in our choice of the most effective treatment immediately post-accident.

Patients and Methods. In this prospective study (2014 – 2016) we treated 29 patients with blow-out fractures. 18 (62%) were treated conservatively and in 11 (38%) we proceeded surgically. We decided whether surgical or non-surgical therapy was appropriate on the basis of clinical ENT, eye examination and the total volume of prolapsed orbital soft tissue. All procedures were performed by the same operating team with a uniform subciliary approach and using PMR splints adapted to the correct size and shape.

Results. On the basis of the mathematical model we reassessed findings in 2 patients: in one we decided against a surgical solution and in the other a surgical approach was indicated. All 18 patients treated conservatively, fully recovered and are free of diplopia. The 11 operated patients are also free of diplopia, only 1 patient (3%) displays clinically insignificant postoperative diplopia in extreme positions when looking upwards.

Conclusion. With proper selection of the optimal treatment, the rate of complete disappearance of diplopia and fully preserved motility of the eyeball ranges from 91 to 97%. Surgical treatment of orbital floor fractures is important, mainly to minimise persistent post-traumatic diplopia which significantly reduces a patient's quality of life.

Key words: blow-out fractures, volumetry, diplopia, CT-assisted surgery

Received: March 24, 2017; Accepted with revision: August 22, 2017; Available online: August 31, 2017
<https://doi.org/10.5507/bp.2017.037>

^aDepartment of ENT and Maxillofacial surgery, 3rd Faculty of Medicine, Charles University in Prague and Military University Hospital Prague, Czech Republic

^bFaculty of Military Health Science in Hradec Kralove, University of Defence, Hradec Kralove, Czech Republic

^cEye Clinic FBMI CVUT Prague, Czech Republic

Corresponding author: Daniel Kovar, e-mail: daniel.kovar.dk@gmail.com

INTRODUCTION

Clinical symptoms of blow-out fractures may be seen in various combinations, but fracture symptoms are sometimes very poorly visible. The presence of an orbital floor fracture, its size and weight, are determined by CT imaging (coronal and sagittal sections) and mathematical calculation of the volume of prolapsed soft orbital tissue.

On the basis of our previous retrospective study, we identified two critical volumes for indicating surgical treatment in blow-out fractures: 500 mm³ in anterior and posterior fractures and 1400 mm³ in anteroposterior fractures. We focused on the creation of new quantitative indication criteria for surgical therapy by calculating the volume of prolapsed soft tissues of the orbit and determining the critical values^{3,4}.

OBJECTIVES

To use a mathematical model to calculate the volume of prolapsed soft tissue of the orbit in blow-out fractures of the orbital floor as an indication criterion for treatment.

In cases of fractures of the orbital floor, to identify immediately, post-trauma patients requiring surgical treatment.

To avoid incorrect treatment indications, for either conservative or surgical therapy.

To prevent or minimise cases of permanent traumatic diplopia as a consequence of inappropriate treatment.

PATIENTS AND METHODS

In our prospective study, we evaluated 29 patients with blow-out fractures of the orbital floor, treated in our department between 2014 and 2016. We treated 11 patients (38%) surgically and in 18 patients (62%) we opted for a conservative approach. The ratio of male to female was 6:1 and of left to right orbit, 4:3. The average age of patients was 39.3 years (19-80 years). Those who were not operated were monitored until the disappearance of the diplopia, while operated patients were monitored on a regular basis: at 1 week, 1 month, 3 months, 6 months and 1 year after surgery. The mechanism of injury most often involved direct blows, followed by falls, sports injuries and car accidents (Table 1).

Table 1. Mechanism of injury.

Mechanism of injury	Blows	Falls	Sports injuries	Car accidents
Rate (%)	52	31	14	3

Previously, we decided on the most appropriate method of treatment in blow-out fractures immediately post-injury on the basis of fracture identification (using CT images of coronary and sagittal planes) and clinical signs (clinically significant and progressive enophthalmos and diplopia, persisting double vision after recovery from edema of the orbital soft tissues and limited mobility of the eyeball).

Since 2014, we have also been able to use the following indication criteria: if, in anterior and posterior fractures, the volume of the prolapsed orbital soft tissue exceeds the critical value of 500 mm³ or 1400 mm³ in anteroposterior fractures.

If no urgent operation is indicated, then the decision about the need to operate or not is made on the basis of the development of clinical symptoms and the total volume of the prolapsed orbital tissue.

All procedures were performed by the same operating team with a uniform subciliary approach and using PMR splints adapted to the correct size and shape.

The main measure of success of the treatment process was the disappearance of diplopia.

RESULTS

To calculate the volume of the prolapsed section of the orbit in a blow-out fracture of the orbital floor, we used the formula for half the volume of a rotating ellipsoid: $V = 2/3 \pi abc$.

The resulting volumes of prolapsed orbital soft tissue in the conservative and surgical treatment methods are shown in Table 2 and Table 3. Clinical symptoms are summarised in Table 4.

In our group of patients with fractures of the orbital floor, one received surgical treatment for a posterior fracture with a prolapse of 435 mm³, diplopia without restricted mobility of the eyeball and with significant hematoma of the eyelids and pneumoorbit. Once the swelling had subsided, a conservative approach was adopted, resulting in a gradual disappearance of clinical symptoms, including diplopia, which resolved within 14 days.

In contrast, one patient initially treated conservatively had gradually receding diplopia but after the repositioning of fractured nasal bones, the diplopia persisted when the patient looked up. There was also a slightly limited range of motion and pain in the lower straight eye muscle (with no muscle entrapment). It was a posterior fracture with a prolapse of 898 mm³. The operation was performed after an interval of 4 months, using the standard subciliary approach, freeing the entire prolapsed volume of the orbit and with orbital floor reconstruction using PMR plate. The patient recovered without complications, and has no diplopia.

DISCUSSION

A number of Czech and foreign authors have addressed the issues of the appropriate timing and method of surgery in blow-out fractures.

Kwon et al. categorise orbital floor fractures as anterior, posterior and anteroposterior⁶ and because of its simplicity and clarity we used the same categorisation in our publications.

Harris, in line with most surgeons^{3,6,9} recommends

Table 2. Prolapsed volumes in blow-out fractures/conservative therapy.

Conservative approach			
Fracture location	Anterior	Posterior	Antero-posterior
Volume (mm ³)	111.6 (36–225)	208.8 (16–435)	0

Table 3. Prolapsed volumes in blow-out fractures/surgical therapy.

Surgical approach			
Fracture location	Anterior	Posterior	Antero-posterior
Volume (mm ³)	856.5 (480–1233)	891.4 (350–1450)	1802.0 (1507–2112)

Table 4. Clinical symptoms.

	Post-traumatic diplopia	Limited eye movement	Exophthalmos	Enophthalmos	Hematoma/pneumoorbit	Post-op diplopia
Operated (%) n=11	91	73	18	9	100	9
Not operated (%) n=18	28	6	11	0	100	0

Table 5. The frequency of blow-out fractures by location and the relationship between conservative and surgical therapy.

Fracture location	Anterior (11), 38%		Posterior (14), 48%		Antero-posterior (4), 14%	
Treatment	Conserv.	Surgical	Conserv.	Surgical	Conserv.	Surgical
No. of fractures	9	2	9	5	0	4
No. of fractures (%)	82	18	64	36	0	100

surgery for defects of the orbital floor where clinically significant enophthalmos can be expected, or in persistent and non-decreasing diplopia 14 days post-accident.

Except in cases of acute danger to the eyeball or if the optic nerve required urgent surgery, procedures were performed 4–5 days after the reduction of edema of the orbital soft tissue, however not more than 14 days after surgery, an approach with which a number of authors are in agreement. This waiting period allows even borderline findings to be monitored.

Again, in various authors, we note the very large dispersion of functional results, 8.7–67%, especially in the persistence of postoperative diplopia^{3,7,10-15}.

Worth noting is the contribution of CT-guided surgery, leading to the complete disappearance of postoperative diplopia. The first functional results with CT navigation were not published until 2014 (ref.^{3,4}).

Beumer, Pham and Schramm¹⁶⁻¹⁸ recommend the use of CT navigation in maxillofacial bone fractures to minimize post-traumatic enophthalmos and achieve improved facial symmetry.

In their conclusion, Mottl et al. emphasise the need for an individual approach to each patient with a fractured orbital floor, stressing this even in non-surgical treatment. The goal of surgical treatment then is the best possible reconstruction of the anatomical shape of the orbit, using new surgical approaches and reconstructive materials¹⁹.

CONCLUSION

Given that on average it is repeatedly seen that less than 50% of patients with orbital floor fractures require surgical treatment, it is important to be able to predict these patients immediately after the injury. Clinical symptoms shortly after injury are affected by hematoma, pneumoorbit and subcutaneous emphysema, which usually change the clinical picture towards the expected pathology (which subsides in a few days), or they appear falsely to be functionally insignificant (symptoms appearing after a period of latency).

In anteroposterior fractures we recommend surgery in prolapse of more than 1400 mm³ and in anterior and posterior fracture types, in cases over 500 mm³ after subsidence of acute edema, provided the situation is non-urgent (Table 5, Fig. 1).

The accuracy of our mathematical model is supported by excellent anatomical and functional results with minimal permanent diplopia, based on experience with CT navigated and video assisted CT-assisted surgery and con-

Blow-out fractures (29), 100%




Fig. 1. Relationship between conservative and surgical therapy and fracture location.

sistent and long-term monitoring of patients, including of outcomes in conservative treatment.

Thanks to the introduction of CT-guided surgery in orbital floor fractures, we are convinced of the importance of freeing the full prolapsed volume of the orbit and completely covering the defect of the orbital floor. With these new methods and resorbable materials, especially thermoplastic and resorbable PMR plate, we are able to achieve excellent reconstructions of the anatomical shape of the orbital floor.

With proper selection of treatment, the rate of complete disappearance of diplopia and fully preserved motility of the eyeball ranges between 91 and 97%.

Surgical treatment of orbital floor fractures is important mainly to minimise persistent post-traumatic diplopia which significantly reduces a patient's quality of life.

Author contributions: DK reviewed the literature, provided the methods and drafted the manuscript. All authors contributed equally to data collection, analysis, interpretation and correction of the final manuscript version.

Conflict of interest statement: The authors state that there are no conflicts of interest regarding the publication of this article.

REFERENCES

- Hoffmann J, Cornelius CP, Groten M, Probst L, Pfannenberger C, Schwenzer N. Orbital reconstruction with individually copymilled ceramic implants. *Plast Reconstr Surg* 1998;101:604-12.
- Galanski M, Friedmann G. Röntgenanatomische Gesichtspunkte zur Diagnostik von Orbitabodenfrakturen. *Fortschr Kiefer Gesichtschir* 1977;22:26-8.

3. Kovář D, Voldřich Z, Voska P, Lešták J, Drahoušková E. Effect of CT Navigation in Surgery of Blow-out Fractures of the Orbital Floor on Functional and Anatomical Results. *Indian Journal of Applied Research* 2014;11:23-7.
4. Kovář D, Voldřich Z, Voska P, Lešták J, Drahoušková E. Video-assisted and CT navigated reposition of an orbital blow-out fracture. *Mil Med Sci Lett (Voj.Zdrav. Listy)* 2014;83:145-50.
5. Kwon JH, Kim JG, Moon JH, Cho JH. Clinical analysis of surgical approaches for orbital floor fractures. *Arch Facial Plast Surg* 2008;10:21-4.
6. Burnstine MA. Clinical recommendations for repair of isolated orbital floor fractures: an evidence-based analysis. *Ophthalmology* 2002;10:1207-13.
7. Cigánek L, Voldřich Z, Lešták J. Diplopia in blow-out fractures. *Čs Oftal* 1990;46:139-44. (In Czech)
8. Harris GJ. Orbital blow-out fractures: surgical timing and technique. *Eye* 2006;20:1207-12.
9. Harris GJ, Garcia GH, Logani SC, Murphy ML, Sheth BP, Seth AK. Orbital blow-out fractures: Correlation of preoperative computed tomography and postoperative ocular motility. *Tr Am Ophth Soc* 1998;XCVI:329-47.
10. Ceylan OM, Uysal Y, Mutlu FM, Tuncer K, Altinsoy HI. Management of diplopia in patients with blowout fractures. *Indian J Ophthalmol* 2011;59:461-4.
11. Gunarajah DR, Samman N. Biomaterials for repair of orbital floor blowout fractures: a systematic review. *J Oral Maxillofac Surg* 2013;71:550-70.
12. Hrušák D, Jambura J, Hauer L. Blow-out Fractures of the Orbital Floor (Statistics) (In Czech). *Čes Stomat* 2010;110:104-8.
13. Chi MJ, Ku M, Shin KH, Baek S. An analysis of 733 surgically treated blowout fractures. *Ophthalmologica* 2010;224:167-75.
14. Jin HR, Lee HS, Yeon JY, Suh MW. Residual diplopia after repair of pure orbital blowout fracture: the importance of extraocular muscle injury. *Am J Rhinol* 2007;21:276-80.
15. Loba P, Kozakiewicz M, Nowakowska O, Omulecki W, Broniarczyk-Loba A. Management of persistent diplopia after surgical repair of orbital fractures. *J AAPOS* 2012;16:548-53.
16. Beumer HW, Puscas L. Computer modeling and navigation in maxillofacial surgery. *Curr Opin Otolaryngol Head Neck Surg* 2009;17:270-3.
17. Pham AM, Rafii AA, Metzger MC, Jamali A, Strong EB. Computer modeling and intraoperative navigation in maxillofacial surgery. *Otolaryngol Head Neck Surg* 2007;137:624-31.
18. Schramm A, Suarez-Cunqueiro MM, Rücker M, Kokemueller H, Bormann KH, Metzger MC, Gellrich NC. Computer-assisted therapy in orbital and mid-facial reconstructions. *Int J Med Robotics Comput Assist Surg* 2009;5:111-24.
19. Mottl R, Slezák R, Feuermannová S, Laco J, Mottlová A, Janovská Z. Isolated Blow-out Fractures of the Orbital Floor. *Praktické zubní lékařství* 2014;62:61-8. (In Czech)