

Radiotherapy dose limit for uterus fertility sparing in curative chemoradiotherapy for rectal cancer

Radka Lohynska^a, Michaela Jirkovska^b, Alena Novakova-Jiresova^a, Eva Mazana^a, Kamil Vambersky^c, Tomas Veselsky^c, Anna Kindlova^c, Hana Stankusova^b, Bela Malinova^b

Aims. Curative sphincter sparing radiotherapy is a treatment option for early rectal cancer. There are many methods developed for fertility preservation in young patients treated with pelvic radiotherapy. Pregnancy rates after radiotherapy are dependent on the radiation dose to ovaries and uterus. Data on outcomes of total body irradiation suggest a pregnancy is possible following 12-14 Gy TBI, despite elevated rates of preterm deliveries and other complications.

Methods. We report a case of full-term delivery of twins after curative chemoradiotherapy for anorectal adenocarcinoma T2 N0 M0 with the total dose 58.6 Gy. The patient underwent laparoscopic laterocranial ovarian transposition before radiotherapy.

Results. Long term complete remission was achieved after treatment. Although a spontaneous conception was not successful, the patient underwent an in vitro fertilisation procedure with donor eggs and conceived twins 10 years after the radiotherapy treatment. The mean dose to the uterus was 16 Gy and to the uterine cervix 35 Gy. She reached a full-term pregnancy and delivered two healthy babies by caesarean section at a gestational age of 38 weeks, weighing 2420 g and 2220 g.

Conclusion. This is the first case report of the successful pregnancy following sphincter sparing curative pelvic radiotherapy for rectal cancer. Furthermore it allows us to propose an increased limit dose to the uterus enabling fertility sparing beyond the limits achieved from total body irradiation series with 12-14 Gy and accept 16 Gy as uterine body (35 Gy for uterine cervix) limit for IMRT treatment planning in young patients asking for maintaining fertility potential.

Key words: rectal adenocarcinoma, radiotherapy, fertility sparing, uterus dose, dose constraints

Received: June 17, 2020; Revised: August 26, 2020; Accepted: September 4, 2020; Available online: September 21, 2020

<https://doi.org/10.5507/bp.2020.039>

© 2021 The Authors; <https://creativecommons.org/licenses/by/4.0/>

^aDepartment of Oncology, First Faculty of Medicine, Charles University in Prague and Thomayer Hospital Prague, Czech Republic

^bDepartment of Oncology, Second Faculty of Medicine, Charles University in Prague and University Hospital Motol Hospital, Prague, Czech Republic

^cDepartment of Medical Physics, University Hospital Motol, Prague, Czech Republic

Corresponding author: Radka Lohynska, e-mail: radka.lohynska@ftn.cz

INTRODUCTION

Radiation therapy in young women has long-term effects on fertility depending on age, radiation dose and site of radiotherapy. Human oocytes are very sensitive to radiation and the estimated median lethal dose (LD50) is less than 2 Gy (ref.¹). The radiation exposure leads to premature ovarian failure. The radiotherapy effect is dependent on patient's age and the mean dose of fractionated radiotherapy to the ovarian tissue and uterus. The effective sterilizing dose (which leads to premature ovarian failure in 97.5% of patients) decreases with increasing age of patient. It is 20.3 Gy at the time of birth, 18.4 Gy at 10 years, 16.5 Gy at 20 years, 14.3 Gy at 30 years and only 6 Gy in women beyond 40 years of age².

Women after pelvic radiation have not only an increased risk of premature ovarian failure, but suffer also from uterine dysfunction associated with miscarriages, placental abnormalities, preterm labour and low birth-weight. Uterine growth is complete around the age of 20 (ref.³). Radiation exposure of the uterus leads to reduced

vascularity, myometrium fibrotic changes and hormone dependent endometrial insufficiency, which reduces subsequent reproductive outcomes due to smaller uterine volumes and atrophic endometrium. In adults, the uterus exposure of 12-14 Gy is associated with significant uterine damage. Childhood exposure to uterine doses of >25 Gy leads to irreversible damage⁴. The highest risk of infertility, miscarriages, preterm labours, intra-uterine growth retardation and low birth weight occurs within one year after radiotherapy⁵. Some authors recommend that patients receiving >45 Gy during adulthood and >25 Gy in childhood should avoid attempting pregnancy⁶.

Clarity on the radiation dose to the uterus, above which a pregnancy would not be sustainable, is lacking. Evidence of successful pregnancies can be found in the literature, for example, a report of IVF assisted conception and full-term delivery 14 years after a high-dose chemotherapy and 54 Gy of pelvic irradiation for Ewing's sarcoma (dose to the uterus was not specified) (ref.⁷). Here, we are the first to report a case of full-term delivery of twins following a curative radiotherapy for anorectal carcinoma.

PATIENT AND METHOD

A 32-years old woman without comorbidities opted for a curative sphincter sparing radiotherapy over radical abdominoperineal resection for distal rectal adenocarcinoma T2 N0 M0. She underwent laparoscopic lateral cranial ovarian transposition 14 days before radiotherapy. Radiotherapy was delivered in prone position with empty bladder using 3D conformal radiotherapy technique. The patient received external beam radiotherapy 48.6 Gy in 27 fractions combined with concurrent continuous 5-fluorouracil chemotherapy combined with 10Gy in 2 fractions interstitial anal canal brachytherapy boost (the primary tumour total dose 58.6 Gy). The mean total dose to uterus was 16 Gy and 35 Gy to the uterine cervix. Written informed consent was obtained from the patient and there were no ethical issues regarding the standard treatment procedure. The dose to transposed ovaries was estimated in the range of 10-12Gy, as the ovaries were not marked with a contrast clip and were indistinguishable from bowel loops. Spontaneous conception was not successful and the patient underwent an *in vitro* fertilisation with donor eggs 10 years later at the age of 42 years resulting in full-term delivery with two healthy babies by caesarean section in gestational age of 38+0 weeks. The babies weighed 2420 g (girl) and 2220 g (boy).

DISCUSSION

The radiation exposure of female reproductive organs is an important toxicity issue as almost 50% of cervical cancer, 10% of anal cancer, 5% of colorectal cancer, 2% of uterine or bladder cancer, and number of soft tissue sarcomas occur in women of childbearing potential. The pelvic radiation is often the treatment of choice in management of such cases⁶. Despite low patient numbers of reported successful cases, the published data on fertility outcomes after pelvic radiation serve as a useful indicator of factors impacting chances of fertility preservation.

There have been several methods of fertility preservation developed for these cases: ovarian transposition, gonadal radiotherapy shielding, embryo and oocyte cryopreservation, cryopreservation of ovarian tissue, donor oocytes and gestational surrogacy or ovarian suppression with GnRH analogues or antagonists.

The uterine radiation doses of less than 4 Gy seem to have a negligible impact on fertility⁸.

The uterine radiation exposure in pre-menarche patients has been more often associated with preterm labour. Thereafter, the radiosensitivity of the uterus further decreases with age⁹. Nevertheless, a dose-effect relation has been reported between an increasing uterine exposure and the risk of low birth-weight, with a spontaneous low birth-weight risk of 7.6% (at a 0 Gy dose level) rising to 25.5% and 36.2% at the exposure ranges of 2.5-5 Gy or > 5 Gy, respectively⁹.

Data on outcomes of adult total body irradiation (TBI) suggest a pregnancy is possible following 12 Gy

TBI, despite elevated rates of preterm deliveries and other complications⁶. Furthermore 14 Gy TBI and bone marrow transplant were noted to lead to ovarian follicular depletion, decreased fertility and impaired uterine growth and blood flow. Pregnancies would terminate in an early pregnancy loss or preterm labour¹⁰. The average uterine volume has been reported to decrease to 40% of the normal adult size regardless of oestrogen substitution¹¹. Another study evaluating fertility 12 years after the bone marrow transplant have demonstrated an increased risk of spontaneous abortion (37% versus 7%) and preterm delivery (63% versus 18%) of low birth weight children in female TBI recipients compared to the chemotherapy-only group¹⁰.

Oocyte donation represents a treatment option for patients with premature ovarian failure. In a case report of three childhood cancer patients after a bone marrow transplant and TBI, the oocyte donation has resulted in 33% full term delivery rate and 33% risk of miscarriage¹².

Whole abdominal and/or pelvic radiotherapy (20–30 Gy) in childhood has been linked to a shorter uterine length, reduced endometrial thickness and a decreased blood flow later in life, associated with a mid-trimester pregnancy loss¹³. Ultrasound measurements revealed that uterine length in chemotherapy group was 7.3 ± 0.6 cm (mean \pm 2 SE) versus 4.1 ± 0.8 cm in the group who had received radiotherapy¹⁴. As for the pelvic irradiation in adolescence or adulthood, the threshold dose for uterine damage remains unknown⁶.

A case of a successful pregnancy has been reported in a 25-year-old woman after receiving 30 Gy of pelvic chemo-radiotherapy for anal cancer¹⁵. Another account has described a successful conception using a donor oocyte program 15 years after irradiation of the right hemi-pelvis for Hodgkin's disease. The pelvic dose was 36 Gy and the age at irradiation was 16 (ref.¹⁶). Following a higher dose yet and an earlier age of exposure (14 y), a patient irradiated for Ewing sarcoma by 54 Gy and 10 Gy to the left and right hemi-pelvis, respectively, was able to conceive spontaneously and carry the pregnancy to full term¹⁷.

CONCLUSION

A fertility-sparing pelvic radiotherapy is a feasible treatment option in rectal cancer. Although a precise dose-effect relationship for uterine functional failure has not been established, data from total body irradiation has proven the dose 14 Gy compatible with pregnancy, some authors point to the similarity of the uterus with other glandular organs (e.g. parotid gland) and recommend to limit the absorbed dose to no more than 20-25 Gy (ref.⁹), but widely accepted uterus dose constraint is not established and published cases of successful pregnancies in rectal cancer radiotherapy with precisely defined dose are missing. In the current report, we have shown that the mean uterine dose of 16 Gy and the mean cervical dose of 35 Gy were compatible with a full-term twin pregnancy and delivery. This mean dose allows accept constraints

to maintain uterine function for pregnancy treated with radiotherapy. The need to address the complexities of fertility-sparing approaches in patients undergoing pelvic radiotherapy underscores the importance of treatment planning in the setting of multidisciplinary oncological teams. Further studies and case reports evaluating uterus dose dependence and associated fertility after pelvic radiotherapy are awaited.

Author contributions: RL: manuscript writing; MJ, EM: manuscript revision, literature search; HS, BM: manuscript final approval; ANJ: written English correction, manuscript final approval; KV, AK, TV: dosimetric data evaluation.

Conflict of interest statement: The authors state that there are no conflicts of interest regarding the publication of this article.

REFERENCES

1. Mahajan N. Fertility preservation in female cancer patients: An overview. *J Hum Reprod Sci* 2015;8(1):3-13. doi: 10.4103/0974-1208.153119
2. Wallace WH, Thomson AB, Saran F, Kelsey TW. Predicting age of ovarian failure after radiation to a field that includes the ovaries. *Int J Radiat Oncol Biol Phys* 2005;62(3):738-44. doi: 10.1016/j.ijrobp.2004.11.038
3. Lushbaugh CC, Casarett GW. The effects of gonadal irradiation in clinical radiation therapy: a review. *Cancer* 1976;37(2 Suppl):1111-25. doi: 10.1002/1097-0142(197602)37:2+<1111:aid-cncr2820370821>3.0.co;2-e
4. Larsen EC, Müller J, Schmiegelow K, Rechnitzer C, Andersen AN. Reduced ovarian function in long-term survivors of radiation- and chemotherapy-treated childhood cancer. *J Clin Endocrinol Metab* 2003;88(11):5307-14. doi: 10.1210/jc.2003-030352
5. Chemaitilly W, Mertens AC, Mitby P, John Whitton, Marilyn Stovall, Yutaka Yasui, Leslie L Robison, Charles A Sklar. Acute ovarian failure in the childhood cancer survivor study. *J Clin Endocrinol Metab* 2006;91(5):1723-8. doi: 10.1210/jc.2006-0020
6. Teh WT, Stern C, Chander S, Hickey M. The impact of uterine radiation on subsequent fertility and pregnancy outcomes. *Biomed Res Int* 2014;2014:482968. doi: 10.1155/2014/482968
7. Rodriguez-Wallberg KA, Karlström PO, Rezapour M, et al. Full-term newborn after repeated ovarian tissue transplants in a patient treated for Ewing sarcoma by sterilizing pelvic irradiation and chemotherapy. *Acta Obstet Gynecol Scand* 2015;94(3):324-8. doi: 10.1111/aogs.12568
8. Sudour H, Chastagner P, Claude L, Emmanuel Desandes, Marc Klein, Christian Carrie, Valerie Bernier. Fertility and pregnancy outcome after abdominal irradiation that included or excluded the pelvis in childhood tumor survivors. *Int J Radiat Oncol Biol Phys* 2010;76(3):867-73. doi: 10.1016/j.ijrobp.2009.04.012
9. Ghadjar P, Budach V, Köhler C, Jantke A, Marnitz S. Modern radiation therapy and potential fertility preservation strategies in patients with cervical cancer undergoing chemoradiation. *Radiat Oncol* 2015;10:50. doi: 10.1186/s13014-015-0353-4
10. Sanders JE, Hawley J, Levy W, Gooley T, Buckner CD, Deeg HJ, Doney K, Storb R, Sullivan K, Witherspoon R, Appelbaum FR. Pregnancies following high-dose cyclophosphamide with or without high-dose busulfan or total-body irradiation and bone marrow transplantation. *Blood* 1996;87(7):3045-52.
11. Holm K, Nysom K, Brocks V, Hertz H, Jacobsen N, Müller J. Ultrasound B-mode changes in the uterus and ovaries and Doppler changes in the uterus after total body irradiation and allogeneic bone marrow transplantation in childhood. *Bone Marrow Transplant* 1999;23(3):259-63. doi: 10.1038/sj.bmt.1701569
12. Larsen EC, Loft A, Holm K, Müller J, Brocks V, Andersen AN. Oocyte donation in women cured of cancer with bone marrow transplantation including total body irradiation in adolescence. *Hum Reprod* 2000;15(7):1505-8. doi: 10.1093/humrep/15.7.1505
13. Critchley HO, Bath LE, Wallace WH. Radiation damage to the uterus - review of the effects of treatment of childhood cancer. *Hum Fertil (Camb)* 2002;5(2):61-6. doi: 10.1080/1464727022000198942
14. Critchley HO, Wallace WH, Shalet SM, Mamtara H, Higginson J, Anderson DC. Abdominal irradiation in childhood; the potential for pregnancy. *Br J Obstet Gynaecol* 1992;99(5):392-4. doi: 10.1111/j.1471-0528.1992.tb13755.x
15. Hürmüz P, Sebag-Montefiore D, Byrne P, Cooper R. Successful spontaneous pregnancy after pelvic chemoradiotherapy for anal cancer. *Clin Oncol (R Coll Radiol)* 2012;24(6):455-7. doi: 10.1016/j.clon.2012.03.006
16. de Menezes E, Tuck SM. Pelvic radiotherapy damage to the endometrium causing morbid adherence of placenta. A new risk factor?. *J Obstet Gynaecol* 2007;27(5):526-7. doi: 10.1080/01443610701535766
17. Bath LE, Tydeman G, Critchley HO, Anderson RA, Baird DT, Wallace WH. Spontaneous conception in a young woman who had ovarian cortical tissue cryopreserved before chemotherapy and radiotherapy for a Ewing's sarcoma of the pelvis: case report. *Hum Reprod* 2004;19(11):2569-72. doi: 10.1093/humrep/deh472